

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/85470/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Willmore, Elaine G. and Smith, Michael ORCID: <https://orcid.org/0000-0002-4199-3315> 2016. Scapular dyskinesia: evolution towards a systems-based approach. *Shoulder & Elbow* 8 (1), pp. 61-70. 10.1177/1758573215618857  
file

Publishers page: <http://dx.doi.org/10.1177/1758573215618857>  
<<http://dx.doi.org/10.1177/1758573215618857>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# **Scapular dyskinesia : evolution towards a systems based approach**

**Willmore E.G & Smith, M.J.**

## **Abstract**

Historically scapular dyskinesia has been used to describe an isolated clinical entity whereby an abnormality in positioning, movement or function of the scapula is present. Based upon this, treatment approaches have focused on addressing local isolated muscle activity. Recently however there has been a progressive move towards viewing the scapula as being part of a wider system of movement that is regulated and controlled by multiple factors including the wider kinetic chain and individual patient centred requirements. We therefore propose a paradigm shift whereby scapular dyskinesia is seen not in isolation but considered within the broader context of patient centred care and an entire neuromuscular system.

## **Introduction**

‘Optimal’ scapular motion is considered to be crucial to the functioning of the shoulder and as such, any alteration in scapular kinematics is believed to be a contributing factor to the development of shoulder pathology.<sup>1</sup>In particular, any variation in the amount of scapular upward rotation that occurs is particularly implicated as a predisposing factor in the development of shoulder symptoms. This is because the scapula must upwardly and externally rotate and posteriorly tilt adequately to prevent the humeral head from compressing and shearing against the under surface of the acromion: - one of the proposed mechanisms for producing the syndrome commonly referred to as sub acromial impingement. Based on this widely

held view, the aim of many shoulder rehabilitation programmes is to correct aberrant, local scapula mechanics.

The notion of scapular dyskinesis for the purposes of this article does not include presentations where there is a defined abnormality of the neuromuscular system – for example neuritis, neuropathy, neuropraxia or other forms of peripheral nerve injury.

The purpose of this article is to summarise current concepts and provide the clinician with not only a foundation of reasoning on which to base clinical decision making but some practical suggestions as to how these could be incorporated and utilised in daily practice.

### **What is normal and does it matter?**

Scapulohumeral Rhythm (SHR) is the co-ordination between humeral and scapular movements and was first described by Codman<sup>2</sup> and subsequently popularised by Inman<sup>3</sup>. X-rays were used to calculate scapula movement of seemingly one subject and the conclusion reached that there was an overall constant scapulohumeral rhythm of 2:1 during shoulder flexion and abduction. Considering a full arc of shoulder movement is 180°, this is a result of 120° of glenohumeral movement and 60° of upward rotation.

With the benefit of time and technology, it is clear that SHR is much more complex than then reported 2:1 ratio. Ratios are characteristically inconsistent suggesting that the relationship is variable and non-linear. In a survey of practicing physiotherapists which asked how reliable and valid they felt that the SHR theory is in predicting

pathological sequences in the shoulder complex Kirby et al<sup>4</sup> reported that 77% held the belief that it was reliable. The study by Kirby et al therefore demonstrates that SHR is a widely held belief and that physiotherapists therefore routinely make clinical decisions based on principles that may lack validity .Upward rotation is of course, just one component of scapula movement with internal/external rotation and antero/posterior tipping occurring simultaneously around three different axis.

If, as the evidence suggests, there is a wide range of physiological normal with high degrees of variability within and between individuals<sup>5,6</sup> and populations<sup>7</sup> benchmarking “normal” against “pathological” poses a considerable challenge. There are also variations of scapular movement within individuals depending on the speed at which movement is performed<sup>8</sup>, the load used<sup>9</sup>, whether movement is concentric or eccentric<sup>10</sup>, performed unilaterally or bilaterally<sup>11</sup> or even the hand dominance of the subject<sup>12</sup>. Against this background, establishing what "abnormal" is in the absence of an accepted norm is a fundamental flaw in the traditional concept.

### **Assessing the “abnormal” scapula – The Traditional Model**

Historically, there have been attempts to establish a threshold for what is and what isn't a dyskinetic scapula such as the scapular lateral slide test<sup>13</sup>. Tests have generally involved a) quantitative measures of abnormality or b) visual observation and a subjective judgement by the clinician as to whether or not it is normal.

Although static and/or dynamic scapular asymmetry is commonly integrated into clinical assessment, the collective body of literature undermines its ubiquity. Both quantitative and qualitative methods are of little clinical value when it is considered that:-

- Postural asymmetries of greater than 1.5cm are commonplace in asymptomatic individuals.<sup>7,14</sup>
- Measurement is subject to measurement error and observer bias<sup>15</sup>
- No difference in scapular dyskinesis between symptomatic and asymptomatic shoulders have been found across several studies<sup>16-18</sup>
- Observed dyskinesis may actually be a strategy to optimise certain individuals or cohorts<sup>19-21</sup>
- Only low – moderate reliability and/or agreements is found between therapists regarding identification or categorisation of dyskinesis<sup>22,23</sup>
- Subjects identified by visual and three dimensional tracking as having dyskinesis are no more likely to report symptoms.<sup>24</sup>
- 

There is also the problem of relevance. If the subject of the measurement (in this case whether or not a scapula is dyskinetic) is a flawed concept then the tests themselves are of limited use.

### **An improbable model?**

As has happened so many times in our profession, the original concept of a pure, de-lineated and possibly dogmatic theory and its associated assessment has, over time and with a growing body of research become questionable. A recent consensus of assessment of visual observation of scapula dyskinesis has further reduced attempts to quantitatively assess scapula dyskinesis to a qualitative yes/no category<sup>25</sup>.

The traditional model of assessment of scapular dyskinesis appears to be becoming more implausible (see figure 1). The evidence challenging the existence of and the assessment for scapular dyskinesis is broad based, but to compound matters, it would appear that attempts to correct identified scapular dyskinesis with rehabilitation are largely unsuccessful. McClure et al <sup>26</sup> found that shoulder impingement patients with supposed scapular dyskinesis, as measured with 3-Dimensional motion sensor equipment, had significant improvements in their pain and function following a 6 week exercise programme. The identified dyskinesis however did not change. Similar results were demonstrated in a more recent study by Struyf et al<sup>27</sup> that improvements in pain and function following rehabilitation were not accompanied by alterations in the so called dyskinetic movement.

**Challenges to the evidence regarding scapula dyskinesia:-**

- In the absence of an identified normal, abnormal is an unknown entity.
- What is perceived as abnormal may in fact be a normal adaptation strategy.
- Tests used to supposedly identified abnormalities cannot be claimed to do so given that they lack construct validity.
  - Measurements are unreliable and prone to measurement error and bias.
- A causal relationship between the existence of scapular dyskinesis and the presence of symptoms cannot be established.

**Figure 1**

The whole concept of scapula dyskinesis as an isolated condition is therefore facing some kind of existential crisis – perhaps it doesn't exist at all!

### **Dissolve or evolve – the theory limps on?**

It would be remiss at this point to advocate the dissolution of all things pertaining to aberrant scapula humeral rhythm and throw the baby out with the kinematic bathwater. Whether it is subjectively labelled as normal or abnormal, what *is* observed is the association and interdependence of the neuromusculoskeletal system around the trunk, shoulder girdle and wider kinetic chain.

The end result should be production of smooth, controlled movement between the humeral and scapula components of movement and this requires significant co-ordination. It is perhaps time to evolve our thinking and move towards a systems based approach.

During the course of movement, muscles almost never work in isolation. Co-ordinated, controlled movement is the outcome of a functioning neuromuscular system. If something goes wrong with the movement, it could be attributed to any part of the system. The complexity of this extraordinary process is reliant on an integrated process of sensorimotor control. Such control requires successful interaction of its component parts such as the motor cortex, thalamic system and cerebellum as well as higher cognitive functions such as perception<sup>41</sup>. The interdependence of all of these systems is underpinned by a vast network of neuronal circuitry, which is far from being understood.

### **Scapula Assessment Revisited. If you can affect it – suspect it.**

The very nature of the complexity and the interdependency of all the systems required for motor control render the existence of a litmus test for the presence of scapular dyskinesis extremely unlikely and the search for one a potentially fruitless task. Over the last decade there has been a gradual move in the assessment of musculoskeletal conditions from the limited value of the traditional orthopaedic test toward a more global, systems based approach. The publication of the shoulder symptom modification procedure (SSMP)<sup>28</sup> signalled a paradigm shift in the way shoulder assessment was conducted. This approach involves subjecting patients with shoulder pain to a series of independent manoeuvres in an attempt to reduce the patient's numerical symptom rating scale (NSRS) by 30%. Such interventions include thoracic (postural) correction, scapula movement facilitation, humeral head procedures and neuromodulation procedures. The muscle contraction element of the humeral head and scapula interventions aim to modify muscle activity around the shoulder with the intention of positively altering the pattern of muscular recruitment of the external rotators, humeral head depressors and/or humeral head adductors. If successful, the facilitation of these muscle groups can be incorporated into early treatment programmes.

Due to its close association with the wider kinetic chain, techniques to reduce the appearance of dyskinesis by incorporating elements chain activity, for example single leg standing and standing on tip toes, have been used with demonstrably good effect<sup>29</sup>. The use of isometric glenohumeral external rotation with or without concurrent shoulder flexion as proposed by the SSMP can be easily applied with substantial affect. The exact mechanism by which these procedures reduce



symptoms is unknown but clinically it appears that by accessing posterior rotator cuff activity via glenohumeral external rotation, a patient with what could be described as visible scapular dyskinesis performs this manoeuvre and, as far as the limitations of visual observation allow, the dyskinesis significantly reduces or disappears altogether. If accompanied by a reduction in pain, this would be a favourable response to the procedure. As mentioned previously studies by Tate et al<sup>24</sup> McClure et al<sup>26</sup> and Struyf et al<sup>27</sup> highlighted patient's outcome in terms of pain and function appears to be unrelated to changes in scapular kinematics. It is therefore not alterations to the appearance of dyskinesis *per se* that is the intended aim of the improvement tests but rather the reduction in reported symptoms that can be achieved with their use.

The fact that such a spectrum of different interventions all have the potential to reduce the dyskinesis introduces another thought dimension. If multiple and varying actions all have the potential to “correct” the scapular dyskinesis then there can be no single mechanism by which said scapular dyskinesis occurred in the first instance. Sub group analysis of which type of patients respond best to which type of intervention would be revealing. However, no such predictive measures have been validated and clinicians do not have the luxury of waiting until they are.

In an attempt to signpost therapists or indeed our surgical colleagues the assessment tools in figure 2 are suggested as a starting place to attempt to find an intervention that affects either the patient's symptoms that may include pain and/or instability.

Figure 2 INSERT CLINICAL APPLICATIONS HERE

### **Making sense of the assessment**

If these 'improvement' or 'symptom modification' procedures have a significant effect on symptoms they could be used at the starting point for rehabilitation and exercise prescription. However, some elements are easier to integrate than others. For example, if dynamic external rotation or tip toe/single leg standing reduces the symptoms or the appearance of dyskinesia then this can easily be used as a foundation on which any number of other exercises functionally relevant to the patient can be built. If however, humeral facilitation or scapular upward rotation improves the patient's symptoms then this is less straightforward. Scapular upward rotation is not a movement that exists in isolation and the translational anterior posterior (AP) glide movement of humeral head facilitation requires an externally applied pressure.

The challenge then becomes finding an exercise that capitalises on the symptom reduction achieved during symptom modification. For this, clinical reasoning needs to take one step further and this is significantly aided by understanding the roles and function of the scapula and rotator cuff.

### **Understanding the functional anatomy**

Type	Role
Agonist	mover – prime or assistant mover
Antagonist	muscle which must relax or work eccentrically to allow the desired movement to occur

Stabiliser	muscle that supports a body part so that another muscle will have a firm base from which to act
Synergist	muscle that eliminates an undesired movement that would otherwise be produced by the mover(s) and stabiliser(s)

Figure 3: The varying roles of muscles

<b>Scapulohumeral Muscles</b> <ul style="list-style-type: none"> <li>• Rotator cuff</li> <li>• Deltoid</li> <li>• Teres major</li> <li>• Long head of triceps</li> </ul>	<b>Axioscapular Muscles</b> <ul style="list-style-type: none"> <li>• Trapezius</li> <li>• Levator scapulae</li> <li>• Rhomboid major + minor</li> <li>• Serratus anterior</li> <li>• Pectoralis minor</li> </ul>	<b>Axiohumeral Muscles</b> <ul style="list-style-type: none"> <li>• Pectoralis major</li> <li>• Latissimus dorsi</li> </ul>
--	--	---

When glenohumeral movement occurs, the scapula must also move to allow the repositioning of the glenoid fossa thus increasing the available range of movement. During this movement, the scapulohumeral and axioscapular muscles must collectively function to maintain optimal mechanical alignment. Rotator cuff activity prevents unwanted humeral head translation but left unchecked, would pull the scapula laterally, essentially creating a destabilising force. The axioscapular muscles respond by preventing the scapulohumeral muscles from destabilising the scapula and produce the upward rotation, posterior tilting and externally rotate necessary for optimal movement and function<sup>30</sup>. A high correlation therefore exists between the action of scapulohumeral and axioscapular muscles and activity in one group does not exist without corresponding activity in the other.


Just like the rotator cuff, scapula muscles have multiple roles which vary, dependent on the task, the load, the speed or the range in which the movement is occurring<sup>30</sup>. At any time, dependent on these factors, any part of the rotator cuff, other scapulohumeral or axioscapular muscles could be acting in an agonist, antagonist, stabilising (static or dynamic) or synergistic function (see figure3). Glenohumeral external rotation exercises for example, are classically regarded as working the rotator cuff in its agonist role. This is true, but it is also true of *all* shoulder rotator torque generators and is therefore not specific to the rotator cuff. Glenohumeral external rotation however *also* requires the scapular muscles to function in their stabilising role, explaining why either static or dynamic rotation utilised through symptom modification procedures can change apparent scapula dyskinesis.

In addition to their agonistic role specific shoulder muscles also stabilise against destabilising forces created by other shoulder muscles but this is far from static. Rotator cuff muscles prevent unwanted translation of the humeral head caused by other humeral muscles, for example deltoid and pectoralis major. Axioscapular muscles work against the destabilising force of the scapulohumeral muscles to prevent the scapulohumeral muscles from translating the scapula off the thoracic cage. To rehabilitate the stabilising function of the axioscapular muscles (and also by virtue of their interdependence - the rotator cuff) it is perhaps more useful to consider strategies that allow the shoulder complex to react to de-stabilising forces. <sup>30</sup>

If a particular movement cannot be performed without apparent scapula dyskinesis, it is worth considering that the cause of this is not “weakness” and inability of, for example serratus anterior to perform its agonistic role but the inability of the scapular


and/or rotator cuff muscles to work simultaneously in both their mobiliser and dynamic stabilising roles. Delayed activation of serratus anterior in the early stages of movement has been implicated as a potential source of scapula dyskinesis in patients with shoulder symptoms.<sup>31,32</sup> Data from these studies has indicated that such movement lag has normalised before 90 degrees which is out with the range of the painful arc classically associated with impingement type symptoms. Although conjecture, there is speculation that any increased rate of upward rotation and/or posterior tipping to overcome latency contributes to the appearance of dyskinesia.

When interpreting assessment findings it is perhaps helpful to consider the following the process outlined in figure 4.




**Fig 4 – Assessment findings and treatment planning**

Given their interdependence, a thorough assessment of the rotator cuff should also be conducted (see figure 5) to complement the scapula assessment.



**Fig 5: Through range assessment of the rotator cuff. The scapula is stabilised therefore not working in its dynamic role**

If weakness of a portion of the rotator cuff is found, together with the findings from symptom modification tests, a clearer picture about where to start with rehabilitation may start to materialise. An example of this process is shown in figure 6.



**Figure 6 Assessment reasoning**



The kinetic chain elements of assessment should not be forgotten. Single leg standing or tip toe standing may not have resulted in the most significant change with symptom modification but they remain a significant part of the bigger picture. If the humeral contingent of the system is unable to perform a pattern of movement without the involvement of the scapular muscles, the scapula is unfairly implicated as the culprit when it is being utilised in an unconventional way in an attempt to maintain function. Similarly, the dyskinesia may not become apparent until the task becomes more complicated and involves co-ordination with the wider chain. Patients may also present differently in an open or closed chain position (see figure 7) In either case, the observed result is faulty and inefficient motor patterns that may be the cause of or equally the response to pain and/or weakness elsewhere in the system.



**Fig 7** The dyskinesia of this breakdancer disappears with closed chain loading

### **Rehabilitation – isolate or integrate?**

The key to success with this approach to rehabilitation is having a sound appreciation of what the scapula and rotator cuff muscle groups are doing with each exercise, why they may or may not be beneficial and how to adapt them if the patient

is unable to perform them without compensatory movements or symptoms. With this knowledge and armed with some clinical reasoning, creativity and often trial and error, it should be possible to find at least one exercise that a patient can leave with, empowered by the knowledge that they can affect their own symptoms with movement.

As previously discussed, it is difficult to justify a position of claiming that a patient is undergoing rotator cuff *or* scapula specific rehabilitation. To suggest that a particular exercise is purely a “scapula stability” exercise may not only be difficult to substantiate based on the research evidence<sup>33</sup> it is also contradictory to the functional anatomy of the region. What one exercise may do more than another however is bias the exercise towards a situation where, for example, the scapula is only working in a static stabilising, rather than a dynamic stabilising role - the former generally being considered an easier or less complex task.

The interdependence of the scapula, rotator cuff and kinetic chain requires clinicians to question the desirability of isolating one part of the system from the other. If the scapular and rotator cuff components are required to work together functionally, they should be trained as such. “Scapula specific rehabilitation” may sound impressive but unless you are engaged in an activity that does not involve any movement of the humerus, it is in all likelihood an unobtainable goal. Furthermore, it is also not possible to isolate single muscles within a single exercises – even activities that demonstrate maximum activation do not do so to the exclusion of other muscles.<sup>34</sup> These lessons have been learned and are well supported with multiple investigations into both exercise and the overwhelming number of supposed specific manual

muscle tests that are proposed for the identification of specific shoulder pathology<sup>35</sup>

As it is almost impossible and generally undesirable to isolate individual muscles, the case for integration strengthens.

### Linking assessment to rehabilitation

The flowchart in figure 6 illustrates how assessment findings may influence the clinicians decision making process. The following exercises are not intended to be prescriptive but to illustrate how different exercises or progressions can influence the relative contributions from the scapula, rotator cuff and kinetic chain.


## **INSERT REHAB SECTION HERE**

### **Strength or motor control?**

Over recent years there have been a wealth of studies utilising electromyographic (EMG) data<sup>36-40</sup> With such a plethora of such information now available in the literature, an unintended consequence of this may be that there has been an unwitting over emphasis on EMG output and equating it with strength. The traditional model of scapular assessment relies on increasing strength by progressing through a rainbow of theraband resistance and inhibitory techniques for supposed “over active” muscles such as taping, trigger point release or massage Exercise progression using a systems based approach would instead focus not on biomechanically correcting scapula position, but on regaining and retraining the whole motor control pattern that had been identified through careful assessment, as being problematic for that patient.



## Scapular Dyskinesis – Traditional Model versus a Symptoms and Systems Based Approach



## **Conclusion**

A functioning proprioceptively mediated motor control system requires co-ordination and therefore integration, of all the different body parts throughout the kinetic chain necessary for that particular movement pattern. For the shoulder this will require assessment of the constituent parts working in both their agonist and stabiliser roles throughout full ranges of both concentric and eccentric phases of movement. It will also necessitate involvement of the wider kinetic chain to train relevant motor patterns that resemble the functional demands of the patient. In the early stages of rehabilitation, it may be necessary to limit the number of simultaneous functions the patient is required to perform. Low load, unsupported shoulder rotation tasks can be used to train complex normal motor patterns therefore the early part of the motor pattern can be recruited and retrained in the appropriate manner. As rehabilitation progresses, the patient is challenged to maintain the correct motor pattern despite the increasing demand and complexity of the relevant task. This may involve elements of speed, co-ordination, load and specific functional requirements relevant to the individual until a normal, fully functioning motor control pattern can be established and crucially, reinforced with repetition. By utilising the fundamental principles of motor learning and skill attainment with what is understood about proprioceptively mediated musculoskeletal rehabilitation we can use our skills to their full potential. The coalescence of assessment and treatment techniques results in a symptoms and systems approach that is patient centred and relevant to the functional requirements of the individual. We advocate that this has advantages over a traditional biomechanical model that utilises assessment principles focused on minute differences in centimetres and degrees and deviations from a normal that arguably doesn't exist.

## References

1. Sauers EL. Clinical evaluation of scapular dysfunction. *Athletic Therapy Today* 2006; 11:10-14.
2. Codman EA. The Shoulder. Rupture of the supraspinatus tendon and the lesions in or about the subacromial bursa. *Privately Printed*. Thomas Dodd Company. Boston
3. Neer CS. Impingement Lesions. *Clinical Orthopaedics and Related Research* 1972; 173: 70-77.
4. Kirby K, Showalter C and Cook C Assessment of the importance of glenohumeral peripheral mechanics by practicing physiotherapists. *Physiotherapy Research International* 2007; 12: 136-146.
5. Nijs J, Roussel N, Vermeulen K, Struyf F, Mottram, S and Meeusen R. Clinical assessment of scapular positioning in patients with shoulder pain: state of the art. *Journal of Manual and Manipulative Therapeutics* 2007; 30:69-75.
6. Morais NV and Pascoal. Scapular positioning assessment: Is side to side comparison clinically acceptable? *Manual Therapy* 2013, 18: 46-53.
7. Oyama S, Myers JB, Wassinger CA, Ricci RD and Lephart SM. Asymmetric resting position in healthy overhead athletes. *Journal of Athletic Training* 2008; 43:565-570.
8. Sugamoto K, Harada, T and Machida A. Scapulohumeral rhythm: relationship between motion, velocity and rhythm. *Clinical Orthopaedics*; 401: 119-124.
9. Forte FC, de Castro MP, de Toledo JM, Ribiero DC and Loss JF. Scapular kinematics and scapulohumeral rhythm during resisted shoulder abduction – Implications for clinical practice. *Physical Therapy in Sport* 2009; 10:105-111.
10. Braman JP, Engel SC, LaPrade, RF, and Ludewig, PM. In vivo assessment of scapulohumeral rhythm during unconstrained overhead reaching in asymptomatic subjects. *Journal of Shoulder and Elbow Surgery* 2009; 18: 960-967.
11. Klopčar N and Lenarčič J. Bilateral and unilateral shoulder girdle kinematics during humeral elevation. *Clinical Biomechanics* 2006. 21:s20-26.
12. Morais NV and Pascoal AG Scapular positioning assessment: Is side to side comparison clinically acceptable? *Manual Therapy* 2013, 18: 46-53.
13. Kibler WB. The role of the scapula in athletic function. *The American Journal of Sports Medicine* 1998; 26:325-337.
14. Koslow PA, Prosser LA, Strony, GA, Suchecki SL and Mattingly GE. Specificity of the lateral scapular slide test: a reliability and validity study. *Physical Therapy* 2003; 81:799-809.
15. Hayes K, Walton, JR, Szomor, ZL and Murrell GAC. Reliability of 5 methods for assessing shoulder range of motion. *Australian Journal of Physiotherapy* 2001; 47: 289-294
16. Nijs J, Roussel N, Vermeulen K, Souvereinys G. Scapular positioning in patients with shoulder pain: a study examining the reliability and clinical importance of 3 clinical tests. *Archives of Physical Medicine and Rehabilitation* 2005; 86 1349-1355
17. Lukasiewicz AC, McClure P, Michener L, Pratt, N and Sennett B. Comparison of 3-Dimensional scapular position and orientation between subjects with and without shoulder impingement. *Journal of Orthopaedics and Sports Physical Therapy* 1999; 29: 574-586

18. Graichen H, Stammberger T, Bonél H, Wiedeman E et al. Three Dimensional analysis of shoulder girdle and supraspinatus motion patterns in patients with impingement syndrome. *Journal of Orthopaedic Research* 2001; 19: 1192-1198
19. Kardua AR, Kerner PJ, Lazarus, MD. Contact forces in the subacromial space: Effects of scapula orientation. *Journal of Shoulder and Elbow surgery* 2005; 14: 393-399
20. Myers JB, Laudner, KG, Pasquale MR, Bradley JP and Lephart SM. Scapular position and orientation in throwing athletes. *The American Journal of Sports Medicine* 2005; 33: 263-271.
21. Laudner KG, Stanek JM, Meister K. Differences in scapular upward rotation between baseball pitchers and position players. *The American Journal of Sports Medicine* 2007; 35:2091-2095
22. Kibler WB, Uhl TL, Jackson WQ, Brooks PV, Zeller B and McMullen J. Qualitative clinical evaluation of scapular dysfunction: A reliability study. *Journal of Shoulder and Elbow Surgery* 2002; 11: 550-556
23. McClure P, Tate AR, Kareha S, Irwin D and Zlupko, E. A clinical method for identifying scapular dyskinesis. Part 1: Reliability. *Journal of Athletic Training* 2009 44:160-164
24. Tate AR, McClure PW, Kareha S et al. A clinical method for identifying scapular dyskinesis. Part 2: Validity. *Journal of Athletic Training* 2009; 44:165-173
25. Uhl TL, Kibler WB, Gecewich B et al. Evaluation of clinical assessment methods for scapular dyskinesis. *Arthroscopy* 2009; 25:1240-1248
26. McClure PW, Bialker, J, Neff N, Williams G and Karduna A. Shoulder function and 3-Dimensional kinematics in people with shoulder impingement before and after a 6-week exercise programme. *Physical Therapy* 2004; 84: 832-848
27. Struyf F, Nijs J, Mollekens S, Jeurissen I, Truijen S, Mottram S, Meeusen R. Scapular-focused treatment in patients with shoulder impingement syndrome: a randomized clinical trial. *Clinical Rheumatology* 2013; 32: 73-85
28. Lewis JS. Rotator cuff tendinopathy / subacromial impingement syndrome: is it time for a new method of assessment? *British Journal of Sports Medicine* 2009; 43: 259-264
29. Le Leu A. The importance of scapula control in postero-inferior glenohumeral joint instability. *ICES Edinburgh* 2010
30. Boettcher CE, Cathers I and Ginn K. The role of shoulder muscles is task specific. *Journal of Science and Medicine in Sport* 2010; 13:651-656
31. Ludewig PM and Cook TM. Alteration in scapula kinematics and associated muscle activity in people with symptoms of shoulder impingement. *Physical Therapy* 2000; 80:276-291
32. Lukasiewicz AC, McClure P, Michener L et al. Comparison of 3-Dimensional scapular position and orientation between subjects with and without shoulder impingement. *Journal of Orthopaedic and Sports Physical Therapy* 1999; 29:574-586
33. Wattanaprakomkul D, Halaki M, Boettcher C, Cathers, I and Ginn, K. The rotator cuff muscles have a direction specific recruitment pattern during flexion and extension exercises. *Journal of Science and Medicine in Sport* 2011; 14:376-382
34. Boettcher C, Ginn K and Cathers I. The “empty can” and “full can” tests do not specifically activate supraspinatus. *Journal of Science and Medicine in Sport*; 2009; 12:435-439



35. Hegedus E, Goode A, Cambell S et al. Physical examination tests of the shoulder:a systematic review with meta analysis of individual tests. *British Journal of Sports Medicine* 2008; 42:80-92
36. Kibler WB, Sciascia AD, Uhl, TL et al. Electromyographic analysis of specific exercises for scapular control in early phases of shoulder rehabilitation. *American Journal of Sports Medicine* 2008; 36:1789-1798
37. Decker MJ, Hintermeister RA, Faber KJ et al. Serratus anterior muscle activity during selected rehabilitation exercises. *American Journal of Sports Medicine* 1999; 27: 784-791
38. Cools AM, Dewitte V, Lanszweert F et al. Rehabilitation of scapula muscle balance: which exercises to prescribe? *American Journal of sports medicine* 2007; 35:1744-1751
39. De Mey K, Cagnie B, Danneels LA et al. Trapezius muscle timing during selected shoulder rehabilitation exercises. *Journal of Orthopaedic and Sports Physical Therapy* 2009 39:743-752
40. Hardwick DH, Beebe JA, McDonnell MK et al. A comparison of serratus anterior muscle activation during a wall slide exercise and other traditional exercises. *Journal of Orthopaedic and Sports Physical Therapy* 2006; 36:903-910
41. R ijezon U, Clark, NC and Treleaven J. Proprioception in musculoskeletal rehabilitation. Part 1: Basic Science and principles of assessment and clinical interventions. *Manual Therapy* 2015; 20: 368-377

